



## Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin  
Acharya N.G. Ranga Agriculture University  
Lam, Guntur, Andhra Pradesh



### Agromet Advisory Bulletin

Date : 12-11-2021

Weather Forecast of CHITTOOR(Andhra Pradesh) Issued On : 2021-11-12(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-11-13	2021-11-14	2021-11-15	2021-11-16	2021-11-17
Rainfall(mm)	12.0	12.0	20.0	24.0	20.0
Tmax(°C)	26.8	28.7	29.4	29.1	28.9
Tmin(°C)	21.3	22.0	22.0	22.5	22.1
RH-I(%)	95	95	95	95	86
RH-II(%)	65	58	59	64	72
Wind Speed(kmph)	9.0	8.0	8.0	7.0	8.0
Wind Direction(Degree)	50	50	40	40	20
Cloud cover(octa)	6	7	8	8	8

#### Weather Summary/Alert:

Partially Cloudy weather with light to moderate rainfall may prevail in the upcoming 5 days with maximum temp of 26-29°C and minimum temperature of 21-22°C.

#### General Advisory:

In view of rain forecast farmers are advised to harvest matured groundnut after attaining dry weather.

#### SMS Advisory:

In view of heavy rains received farmers are advised to drain out the excess water and protect the crops from fungal diseases.

#### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	• In view of rainfall forecast farmers are advised to provide drainage channels in paddy nurseries to protect the seedlings. After removing excess water apply 15 kg urea and 15 kg potash per acre. • In areas where paddy is at maturity to harvest stage farmers are advised to postpone the harvesting and where the crop is harvested protect the heaps with plastic sheets. • In view of heavy rainfall forecast there is a chance of occurrence of BPH after cessation of rains. To control BPH remove excess water and spray buprofigin @ 1.6 ml./L of water.
BLACK GRAM	In view of rain forecast, In areas of excess rainfall Blackgram farmers are advised to drain the excess water. After cessation of rains, a prophylactic spray with Mancozeb@2.5g/lit of water is recommended to reduce the incidence of diseases.
GROUNDNUT	• In view of rains received farmers are advised to harvest the matured groundnut before germination. • In view of rains received farmers are advised not to take up rabi sowings in excess moisture and advised to take up rabi crops when suitable weather conditions are prevailed.

#### Horticulture Specific Advisory:

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
TOMATO	To control Early blight disease in tomato spray mancozeb or chlorothalonil @ 2 g. or propiconazole 1 ml./L of water twice or thrice at 15 days interval.

**Others (Soil / Land Preparation) Specific Advisory:**

<b>Others (Soil / Land Preparation)(Varieties)</b>	<b>Others (Soil / Land Preparation) Specific Advisory</b>
GENERAL ADVICE	Remove the excess stagnated water from nurseries, just transplanted paddy and crop at tillering wherever excess rains received.