2/28/22, 1:28 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Acharya N.G. Ranga Agriculture University



Lam, Guntur, Andhra Pradesh

Agromet Advisory Bulletin

Date: 28-02-2022

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2022-02-28(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-03-01	2022-03-02	2022-03-03	2022-03-04	2022-03-05
Rainfall(mm)	0.0	0.0	0.0	5.0	18.0
Tmax(°C)	34.6	34.2	34.7	34.5	34.0
Tmin(°C)	17.8	18.2	18.8	19.3	19.5
RH-I(%)	73	75	82	84	88
RH-II(%)	44	45	42	45	48
Wind Speed(kmph)	4.0	5.0	5.0	6.0	4.0
Wind Direction(Degree)	110	90	70	70	70
Cloud Cover(Octa)	2	7	7	7	8

Weather Summary/Alert:

Partially cloudy weather with light rainfall may prevail in the upcoming 5 days with maximum temp of 34-35°C and minimum temperature of 17-19°C.

General Advisory:

The present weather conditions are favourable for leaf hoppers in mango. To control spray fipronil @ 2 g. (or) dimethoate @ 2 ml. (or) imidachloprid @ 0.3 ml. /L of water.

SMS Advisory:

To control spodoptera in groundnut spray thiodicarb @ 1.5-2 g./L of water at evening hours.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
GROUNDNUT	The present weather conditions are favourable spodoptera in groundnut. To control spray thiodicarb @ 1.5-2 g./L of water at evening hours.
RICE	• The present weather conditions are favourable for blue beetle in paddy. To control spray chloripyriphos @ 2.5 ml./L of water. • The present weather conditions are favourable for stem borer in paddy. To control spray cartaphydrochloride @ 2 g./L of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
	The present weather conditions are favourable for leaf hoppers in mango. To control spray fipronil @ 2 g. (or) dimethoate @ 2 ml. (or) imidachloprid @ 0.3 ml. /L of water.

2/28/22, 1:28 PM District Advisory

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The present weather conditions are favourable for pin worm in tomato. To control arrange pheromeone traps @ 10/acre. If the larval attack is more spray korajen @ 0.4 ml./L of water.
	The present weather conditions are favourable forupward and downward leaf curl in chillies. To control spray Fsalon 3 ml (or) Difenthioron 1 g/L of water.