



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Acharya N.G. Ranga Agriculture University
 Lam, Guntur, Andhra Pradesh



Agromet Advisory Bulletin

Date : 06-05-2022

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2022-05-06(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-05-07	2022-05-08	2022-05-09	2022-05-10	2022-05-11
Rainfall(mm)	4.0	0.0	0.0	10.0	0.0
Tmax(°C)	38.5	38.6	38.8	37.1	36.1
Tmin(°C)	25.6	25.8	25.8	24.9	21.8
RH-I(%)	67	66	65	69	72
RH-II(%)	42	40	41	38	46
Wind Speed(kmph)	5.0	5.0	5.0	5.0	5.0
Wind Direction(Degree)	160	160	150	170	160
Cloud Cover(Octa)	6	6	8	8	8

Weather Summary/Alert:

Partially cloudy weather with light to moderate rainfall may prevail in the upcoming 5 days with maximum temp of 37-39°C and minimum temperature of 25-26°C.

General Advisory:

In view of continuous high temperatures farmers are advised to provide drinking water to poultry and cattle at frequent intervals.

SMS Advisory:

Due to continuous high temperatures rainfall is expected at isolated places. Farmers are advised to harvest matured fruits and vegetables and protect the produce on threshing floors.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
GROUNDNUT	• Farmers who are taking up groundnut sowing under irrigation are advised to take up seed treatment compulsory. To control sucking pests treat the seed with Imidachloprid 600 F.S @ 2 ml. + 4 ml water per kg of seed.
RICE	• The present weather conditions are favourable for stem rot in paddy. To control spray tilt @ 1 ml (or) tebuconazole @ 1.5 ml./L of water.
FODDER MAIZE	• Cultivation of Fodder jowar, maize and bajra where irrigation water is available and should harvest after 45-50 days for fodder to cattle.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

Horticulture(Varieties)	Horticulture Specific Advisory
MANGO	• The present weather conditions are highly favorable for occurrence of fruit fly in mango orchards. To control erect pheromon traps or IIHR traps @ 8-10/acre at 3-6 feet from ground level.
TOMATO	The present weather conditions are favourable for pin worm in tomato. To control arrange pheromeone traps @ 10/acre. If the larval attack is more spray korajen @ 0.4 ml./L of water.