



Agromet Advisory Bulletin

Date : 29-07-2022

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2022-07-29(Valid Till 08:30 IST of the next 5 days)

| Parameter | 2022-07-30 | 2022-07-31 | 2022-08-01 | 2022-08-02 | 2022-08-03 |
|------------------------|------------|------------|------------|------------|------------|
| Rainfall(mm) | 6.0 | 12.0 | 42.0 | 30.0 | 50.0 |
| Tmax(°C) | 34.2 | 34.0 | 29.6 | 31.0 | 29.2 |
| Tmin(°C) | 24.5 | 24.0 | 22.5 | 23.0 | 21.8 |
| RH-I(%) | 72 | 76 | 90 | 88 | 94 |
| RH-II(%) | 58 | 60 | 74 | 66 | 80 |
| Wind Speed(kmph) | 8.0 | 10.0 | 9.0 | 10.0 | 12.0 |
| Wind Direction(Degree) | 170 | 220 | 280 | 290 | 290 |
| Cloud Cover(Octa) | 5 | 7 | 8 | 8 | 8 |

Weather Summary/Alert:

Cloudy weather with light to moderate rainfall may prevail in the upcoming 5 days with maximum temp of 30-34°C and minimum temperature of 22-24°C.

General Advisory:

Farmers who are taking up groundnut sowings are advised to raise 5 to 6 row of jowar or bajra as border crops to protect the crop from pests and viral diseases.

SMS Advisory:

The present weather conditions are favourable for fall army worm in maize. To control spray of neem oil @ 5 ml. /L of water

Crop Specific Advisory:

| Crop(Varieties) | Crop Specific Advisory | | | | |
|-----------------|--|--|--|--|--|
| | • The farmers who are taking up transplanting of paddy are advised to appy pre emergence herbicides like pretilachlor @ 500 ml. /acre or oxadiargyl @ 35-50 g. /acre at 3-5 days after transplating by mixing in 20 kg sand/acre keeping thin film of water in the field. • Leave alleyways of 20 cm width for every 2 meters of transplanted crop. • Cut the tip of the seedlings before transplanting. | | | | |
| | • Farmers who are taking up Groundnut sowings are advised to follow seed treatment with tebuconajole @ 1 g/kg of seed. • For the groundnut crop which is in at flowering stage, farmers are advised to apply zypsum @ 200 kg/acre at 5 cm depth followed by intercultivation. | | | | |
| GROUNDNUT | • The present weather conditions are favourable for early leaf spot in early sown groundnut and root rot in Groundnut crop. To control early leaf spot spray hexaconazole @ 2 ml/L of water. To control root rot in 15 days aged crop apply the same fungicide in drenching method. | | | | |

| Crop(Varieties) | Crop Specific Advisory | | | | |
|-----------------|--|--|--|--|--|
| MAIZE | • The present weather conditions are favourable for fall army worm in early vegetative stages in maize. To control spray NSKE @ 5% or arrange pheromone traps @ 10-15 per acre. • If the infestation is high spray spray larvin @ 1 g./L of water. | | | | |

Horticulture Specific Advisory:

| Horticulture(Varieties) | Varieties) Horticulture Specific Advisory | | | |
|-------------------------|---|--|--|--|
| | The present weather conditions are favourable for wilt in tomato. To control spray mancozeb @ 3g/L of water | | | |