

Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Acharya N.G. Ranga Agriculture University Lam, Guntur, Andhra Pradesh



# **Agromet Advisory Bulletin**

Date : 17-03-2023

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2023-03-17(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-03-18	2023-03-19	2023-03-20	2023-03-21	2023-03-22
Rainfall(mm)	35.0	12.0	18.0	12.0	12.0
Tmax(°C)	31.6	32.0	32.8	34.2	34.9
Tmin(°C)	20.6	19.8	21.2	22.4	23.0
RH-I(%)	78	84	81	74	71
RH-II(%)	62	65	56	57	51
Wind Speed(kmph)	8.0	9.0	8.0	9.0	8.0
Wind Direction(Degree)	150	180	180	200	180
Cloud Cover(Octa)	8	7	6	7	2

### Weather Summary/Alert:

Cloudy weather with light to moderate rainfall may prevail in the upcoming 5 days with maximum temp of 31-35°C and minimum temperature of 19-23°C.

#### **General Advisory:**

In view of rainfall forecast and moderate wind speeds, farmers are advised to spray pesticides and fungicides on rain free days in the morning hours.

#### **SMS Advisory:**

In view of rainfall forecast farmers are advised to protect the harvested produce by covering with polythene sheets. In view of moderate wind speeds farmers are advised to protect the crops.

#### **Crop Specific Advisory:**

<b>Crop(Varieties)</b>	Crop Specific Advisory			
GROUNDNUT	• The present weather conditions are favourable for sucking pests in groundnut. To control spray monocrotophos @ 1.6 ml. (or) imidachloprid @ 0.3 ml. (or) acephate @ 1.5 g./L of water.			
RICE	• The present weather conditions are favourable for stem borer in paddy. To control spray chloripyriphos @ 2.5 ml. (or) chloranthraniliprole @ 0.3 ml. /L of water. • The present weather conditions are favourable for BPH in paddy. To control drain the field and spray buprofigin @ 1.6 ml./L or ithophenprox @ 2 ml./L of water.			

District Advisory

Crop(Varieties)	Crop Specific Advisory
SUGARCANE	The present weather conditions are favaourable for stem borer in sugarcane. To control • Remove the late tillers and make the crop stand and provide drainage water facility. • Release trycogama cords @ 20,000/acre from 120 days for every 6 to 8 times at an interval of 1 week to 10 days. • Remove the lower dried leaves and spray chloripyriphos @ 2.5 ml. or monocrotophos 1.6 ml./l of water. • The present weather conditions are favourable for the incidence of early shoot borer in sugarcane. To control go for frequent irrigation at regular intervals and spray chloripyriphos @ 2.5 ml./L of water.

## Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory			
CHILLI	• The present weather conditions are favourable for Upward and downward leaf curl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1 g./L of water.			
MANGO	• The present weather conditions are favourable for hoppers in mango. To control spray imidachloprid @ 0.3 ml. /L of water. • Due to increasing temperatures there is possibility of increased occurrence of thrips in mango. To control spray azadiractin @ 1500 ppm @ 2 ml. or neem oil @ 10 ml./L of water. • Irrigation should be given 2 times in a fortnight. This will help in increasing the fruit size and prevents the fruit fall. • The present weather conditions are favourable for early fruit drop in mango. If the effect is more spray planofix @1 ml./4.5L of water.			
MUSKMELON	The present weather conditions are favorable for thrips in melons which cause vira diseases. To control the vector spray spinosad $@ 0.35 \text{ ml./L}$ of water.			