

Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Acharya N.G. Ranga Agriculture University Lam, Guntur, Andhra Pradesh



Agromet Advisory Bulletin

Date : 20-06-2023

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2023-06-20(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-06-21	2023-06-22	2023-06-23	2023-06-24	2023-06-25
Rainfall(mm)	12.0	8.0	6.0	2.0	2.0
Tmax(°C)	34.8	34.4	33.8	32.2	33.0
Tmin(°C)	25.8	26.2	26.0	25.6	25.4
RH-I(%)	78	80	77	82	85
RH-II(%)	64	62	68	66	70
Wind Speed(kmph)	8	7	8	7	9
Wind Direction(Degree)	290	250	270	270	260
Cloud Cover(Octa)	8	7	6	4	4

Weather Summary/Alert:

Cloudy weather with light to moderate rainfall may prevail in the upcoming 5 days with maximum temp of 32-35°C and minimum temperature of 25-26°C.

General Advisory:

The present weather conditions are congenial for Leaf folder attack in paddy which is at nursery to tillering stage. To control spray cartaphydrochloride @ 2 g/L of water.

SMS Advisory:

In view of rain forecast farmers are advised to go deep ploughings and prepare the land for kharif sowings.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory			
RICE	Farmers who are taking up transplanting of paddy nurseries are advised to • Leave 20 cm pathways for every 2 mt. of transplanted crop • Apply entire 'P2O5' & 'K2O' as basal and 1/3 'N' as basal during lost puddling. • Transplant 30-35 days old rice seedlings. • Maintain the population @ 33 hills/sq.meter			
GROUNDNUT	 Farmers who are taking up Kharif groundnut should select varieties like Narayani, Dharani, K-9, ICGV-9114 and varieties like Grishma, Dheeraj, Nityaharitha, K-6 where availability of water is there and should go for seed treatment with Mancozeb or Tebukonazole @ 3 g/kg of seed. Plant population should be maintained at 33 plants/meter Farmers are advised to clean the bunds and remove the weeds in and around the fields so as to avoid harbouring of pests 			
FINGER MILLET	• Farmers who wish to takeup rain fed small millets are advised takeup sowings from june 2nd fortnight to july 1st fortnight. • Soak the seed for 10-12 hours, shade dry and broadcast the seed in the field at optimum moisture.			

Horticulture Specific Advisory:

District Advisory

Horticulture(Varieties)	Horticulture Specific Advisory			
MANGO	The present weather conditions are highly favorable for occurrence of fruit fly in mango orchards. To control fruit fly erect pheramon traps or IIHR traps @ 8-10/acre at 3-6 feet from ground level (or) spary Spinosad 0.3 ml/L of water.			
• The present weather conditions are favourable for mealy bug in vegeta control spray malathaion or dimethioate @ 2ml and apply with sandwit				