08/08/2023, 14:52 District Advisory



## Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Acharya N.G. Ranga Agriculture University



Lam, Guntur, Andhra Pradesh

# **Agromet Advisory Bulletin**

Date: 08-08-2023

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On: 2023-08-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-08-09	2023-08-10	2023-08-11	2023-08-12	2023-08-13
Rainfall(mm)	1.0	2.0	2.0	1.0	0.0
Tmax(°C)	35.2	34.8	34.5	34.9	34.3
Tmin(°C)	24.5	24.7	24.3	24.8	24.1
RH-I(%)	74	75	73	71	68
RH-II(%)	53	51	52	49	47
Wind Speed(kmph)	6	8	6	8	10
Wind Direction(Degree)	270	280	270	280	280
Cloud Cover(Octa)	5	4	5	5	4

#### **Weather Summary/Alert:**

Partially cloudy weather with very light rainfall may prevail in the upcoming 5 days with maximum temp of 34-35°C and minimum temperature of 24-25°C.

#### **General Advisory:**

Farmers are advised to protect groundnut crop at flowering by providing one irrigation wherever possibility exists.

### **SMS Advisory:**

To control sucking pests in groundnut spray imidachloprid @ 0.3 ml. (or) thiochloprid @ 0.25 ml./L of water.

#### **Crop Specific Advisory:**

Crop(Varieties)	Crop Specific Advisory
RICE	• The present weather conditions are favourable for leaf folder and stem borer attack in tillering stage paddy. To control leaf folder and stem borer attack in paddy spray cartaphydrochloride @ 2 g. /L of water.
GROUNDNIJT	• Rainfed groundnut is at vegetative stage to pegging stage. Due to continuous dry spell and high temperatures the crop is wilting. • Farmers who are taking up intercultivation by providing sprinkler irrigation are advised to spread the removed weeds on the furrows to avoid moisture loss through evaporation.
	• In sugarcane crop there is possibility of incidence of early shoot borer. To control go for frequent irrigation at regular intervals and spray Endosulfan @ 2.0 ml./L. of water.

#### **Horticulture Specific Advisory:**

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The present weather conditions are favourable for pin worm in tomato. To control
	arrange pheromeone traps @ 10/acre. If the larval attack is more spray korajen @

08/08/2023, 14:52 District Advisory

Horticulture(Varieties)	Horticulture Specific Advisory		
	0.4 ml./L of water.		
BRINJAL	The present weather conditions are favourable for stemand fruit borer in brinjal and bhendi. To control spray quinolphos @2ml or monocrotophas 1.6ml or chloripyriphos @ 2 ml./L of water.		