



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Acharya N.G. Ranga Agriculture University
 Lam, Guntur, Andhra Pradesh



Agromet Advisory Bulletin

Date : 05-09-2023

Weather Forecast of District CHITTOOR(Andhra Pradesh) Issued On : 2023-09-05(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-09-06	2023-09-07	2023-09-08	2023-09-09	2023-09-10
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	29.0	31.3	32.4	32.8	33.6
Tmin(°C)	23.7	23.5	24.2	23.8	24.5
RH-I(%)	80	82	79	81	80
RH-II(%)	69	67	65	72	70
Wind Speed(kmph)	6	6	6	6	6
Wind Direction(Degree)	280	250	250	250	250
Cloud Cover(Octa)	4	4	5	5	3

Weather Summary/Alert:

Partially cloudy weather may prevail in the upcoming 5 days with maximum temp of 29-34°C and minimum temperature of 23-24°C.

General Advisory:

Farmers are advised to take up sowings of crop like redgram, castor, greengram, cowpea, sorghum, bajra and horsegram when there is enough moisture in the soil.

SMS Advisory:

To control early leaf spot in groundnut spray spray saaf (bavistin + mancozeb) 2g/L of water.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	• Farmers who are taking up paddy transplanting should apply carbofuran granules @160g/cent to control leaf folder attack in main field. To control BPH leave alleyways of 20cm width for every 2 mt. • The present weather conditions are favourable for stem borer and leaf folder in paddy. To control spray Chloripyrifhos @ 2.5 ml (or) Cartaphydrochloride @ 2 g/L of water.
GROUNDNUT	• The present weather conditions are favourable for Leaf spot in groundnut. To control spray saaf (bavistin + mancozeb) 2g/L of water.
HORSE GRAM	September month is suitable for taking up the sowings of Horse gram. Farmers who are taking up Horse gram are advised to select varieties like PDM-1, VZM-1, PHG-9 and PHG-62 and follow seed treatment before sowing. Inter row spacing should be 30cm and intra row spacing should be 10cm.
PIGEON PEA (RED GRAM/ARHAR)	• Farmers having irrigation facility are advised to take up rabi redgram sowing selectibselecting suitable varieties by maintaining spacing of 60-90 cm between row to row and 10 cm between the plants with a seed rate of 10-12 kg/acre.

Crop(Varieties)	Crop Specific Advisory
MAIZE	<ul style="list-style-type: none"> • The present weather conditions are favourable for fall army worm in Maize. To control farmers should arrange 10 pheromone traps per acre to monitor fall army worm. • For each catch if population exceeds 10, at vegetative stage of the crop to control 1st and 2nd instar larvae spray neem oil @ 5 ml./L of water.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
TOMATO	The present weather conditions are favourable for wilt in Tomato. To control spray mancozeb @ 3g/lit of water.
BRINJAL	The present weather conditions are favourable for stem and fruit borer in brinjal and bhendi. To control spray quinolphos @ 2ml or monocrotophos 1.6ml or chloripyriphos @ 2 ml./L of water.