# Custom Advisory At District Level of Andhra Pradesh:27-02-2024

DISTRICT-CHITTOOR

Weather Forecast of CHITTOOR District in Andhra Pradesh:-Issued On:2024-02-27 (Valid Till 08:30 IST of the

next 5 days)

Date (yyyy-mm- dd)	Rainfall (mm)	Temp Max (°C)	Temp Min (°C)	Humidity I (%)	Humidity II (%)	Wind Speed (kmph)	Wind Direction (Degree)	cloud cover (Octa)
2024-02-28	0.0	33.0	21.0	85	45	6	50	4
2024-02-29	0.0	33.4	21.6	82	48	7	50	4
2024-03-01	0.0	33.7	22.3	80	42	6	200	4
2024-03-02	0.0	33.3	22.0	77	40	7	180	5
2024-03-03	0.0	34.4	22.5	79	43	9	140	4

#### **WEATHER SUMMARY ALERT**

Dry weather may prevail in the upcoming 5 days with maximum temp of 33-34°C and minimum temperature of 21-22°C.

#### **GENERAL ADVISORY**

The present weather conditions are favourable for sucking pests in groundnut. To control spray monocrotophos @ 1.6 ml. (or) imidachloprid @ 0.3 ml. (or) acephate @ 1.5 g./L of water.

# **SMS ADVISORY**

To control pin worm in tomato arrange pheromone traps @ 10/acre. If the larval attack is more spray korajen @ 0.4 ml./L of water.

#### **CROP SPECIFIC ADVISORY**

Select category: crop Select Type: G.Nut Stage: Pod development

• The present weather conditions are favourable for late leaf spot in groundnut. To control spray propiconazol 1 ml./L (or) hexaconazole @ 2 ml./L of water.

Select category: Select Type: Paddy Stage: Vegetative

• The present weather conditions are favourable for stem borer in paddy. To control spray cartaphydrochloride @ 2 g (or) acephate @ 1.5 ml./L of water

### **Select category:**

Select Type: Chilli

Select variety: vegetative

• The present weather conditions are favourable for Upward and downward leaf curl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1 g./L of water.

# Select category:Crop

Select Type:Maize

Select variety: vegetative stage

- The present weather conditions are favourable for fall army worm in Maize. To control farmers should arrange 10 pheromone traps per acre to monitor fall army worm.
- In early stage crop, to control the larvae spray Emamectin benzoate @0.4 g/L. In 35 days crop spray Chloranthraniliprol @ 0.3 ml/L and in matured crop spray Spinecoram 0.5 ml/L of water.

### Select category:

Select Type:

MANGO

Select variety: flowering

- The present weather conditions are favourable for hoppers and anthracnose in mango. To control spray imidachloprid @ 0.3 ml. + carbendazem @ 1 g. (or) thiophenate methyl @ 1 g./L of water.
- The present weather conditions are favourable for early fruit drop in mango. If the effect is more spray planofix @1 ml./L of water.

# **Select category:**

Select Type:

Gingelly

stage: vegetative

• The present weather conditions are favourable for sucking pests in gingelly. To control spray monocrotophos @ 1.6ml. (or) quinolphos @ 2 ml./L of water.

# **Select category: Horticulture**

Select Type: Brinjal

Select variety: fuiting

• The present weather conditions are favourable for fruit borer in vegetables. To control spray chloripyriphos @ 2.5 ml./L of water.